

FOR ANIMAL TREATMENT ONLY

PLEvac[®]

**Parvovirus/Leptospirosis/
Erysipelas Vaccine for Pigs**



ACTIVE CONSTITUENTS:

Inactivated adjuvanted vaccine containing:

Erysipelothrix rhusiopathiae 1 x 10⁹

Erysipelothrix rhusiopathiae 5 x 10⁸

Leptospira interrogans pomona 3.75 x 10⁸

Porcine parvovirus 512 HA/mL

Preservatives: 0.1 mg/mL Thiomersal; < 1.2 mg/mL Formaldehyde

For protection against Erysipelas and *Leptospira pomona* infections and as an aid in the control of parvovirus infection in pigs.

DESCRIPTION:

PLEvac is a vaccine comprised of three antigenic components, Porcine parvovirus, *Erysipelothrix rhusiopathiae* and *Leptospira interrogans* serovar *pomona*. The adjuvant system increases the level of specific antibodies which act as an aid in controlling porcine parvovirus, and for the protection against erysipelas and leptospiral infections in sows and gilts.

SAFETY:

PLEvac has been shown to be safe when administered to pregnant sows at the recommended dose rate.

DIRECTIONS FOR USE:

DOSAGE:

4 mL by intramuscular injection. The recommended site of injection is that used for routine injection in pigs i.e. behind and below the ear.

Gilts are vaccinated at selection then again 3-4 weeks later.

Sows (previously vaccinated) - Vaccinate at 3 weeks before farrowing or at weaning.

CAUTION: Avoid Carcass Damage

1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. Use needles of appropriate gauge and length.
5. As far as possible avoid injection of animals during wet weather or under dusty conditions.

6. This product must be injected only into muscle tissue.
7. If possible inject into muscle tissue on side of neck.

WITHHOLDING PERIOD: NIL

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): ESI Zero (0) days.

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131 126.

CONNECTING PACK TO AUTOMATIC VACCINATOR:

Connect the tube firmly onto the automatic vaccinator. The pack may be suspended by means of a carrying strap. Eject air from the vaccinator and tube by depressing plunger until vaccine flows through needle.

RESEALING VACCINE PACK:

For best practice vaccine should be used during one administration. Unused vaccine must be kept refrigerated and used within 24 hours of opening. Remove tube from the vaccinator gun. Store between 2°C and 8°C (Refrigerate. Do not freeze). Protect from light. **Discard if previously frozen.**

ADDITIONAL SAFETY INFORMATION:

Wash and disinfect self injection injuries. Seek medical advice if irritancy or an allergic response occurs.

DISPOSAL:

Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE:

Store between 2°C and 8°C. (Refrigerate. Do not freeze). Protect from light.

Discard if previously frozen.

APVMA Approval No.: 46164/50907




MSD ANIMAL HEALTH
Intervet Australia Pty Ltd
91-105 Harpin Street,
BENDIGO EAST, VIC 3550
Ph: 1800 033 461

Client: MSD Animal Health

Product Description: PLEvac common Leaflet

Scale: 100%

Colour:  100% Black

Keyline:  (not to be printed)

Dimensions: 80 mm x 150 mm

Clients Code: 144774

PPC: 13-048

Agency Code & Version: MSD14879 V3

Last revised: 9/1/14

Signal Heading: Lucida Sans AK Bold 3 mm

Product Name: Lucida Sans AK Bold 6.5 mm

Body Text: Lucida Sans AK Roman 1.6 mm

Directions for Use: Lucida Sans AK Bold 1.6 mm

Safety Directions: Lucida Sans AK Bold 1.6 mm