

APPROVED

1 x 200 mL (100 doses)

Porcilis[®]
PCV Vaccine
FOR ANIMAL TREATMENT ONLY

127116 R5

DRUG INTERACTIONS:

Compatibility studies with concurrent use of other veterinary products were not performed.

WITHHOLDING PERIOD: Zero (0) days.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): ESI not required.

FIRST AID:

If poisoning occurs contact a doctor or Poisons Information Centre.
Phone Australia 13 11 26.

ADDITIONAL USER SAFETY: Take care to avoid self-injection

This product contains mineral oil. In the event of self-administration, seek prompt medical attention and take this package leaflet/carton with you. Accidental self-administration may result in local bruising, severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If pain persists for more than 12 hours after medical examination, seek medical advice again.

To the physician:

This product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert, PROMPT, surgical attention is required and may necessitate early incision of the injected area, especially where there is involvement of finger pulp or tendon. Additional information is listed in the Material Safety Data Sheet.

Dispose of empty container by wrapping with paper and putting in garbage.

Store between 2°C and 8°C (Refrigerate. Do not freeze). Protect from light.

FOR BATCH AND EXPIRY SEE BOTTOM PANEL

APVMA Approval No.: 64025/47209

Intervet Australia Pty Limited (MSD Animal Health)
91-105 Harpin Street, Bendigo East VIC 3550
Phone: 1800 033 461

® Registered Trademark



9 322839 022765 >



PSA020236 01
65x65x114,6mm
CMYK

FOR ANIMAL TREATMENT ONLY

Porcilis[®]
PCV Vaccine



Porcine Circovirus Type 2 ORF2 subunit antigen:
≥ 4.5 log₂ antibody titre per dose of 2 mL.

Adjuvant: 25 mg dL-α-tocopheryl acetate
346 mg light liquid paraffin

For active immunisation of pigs to reduce the virus load in blood and lymphoid tissues and to reduce mortality and weight loss associated with Porcine circovirus type 2 related infection (PCVD) occurring during the fattening period.

1 x 200 mL (100 doses)



READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

DIRECTIONS FOR USE: SHAKE WELL BEFORE USE. Use entire contents within 8 hours of opening.

RESTRAINTS:

DO NOT mix with any other vaccine or immunological product.

DOSAGE AND ADMINISTRATION:

The vaccine is a liquid and ready-to-use product. Before using the vaccine allow it to reach room temperature (15-25°C) and shake well before use. Use sterile syringes and needles. Avoid introduction of contamination. Avoid use of vaccination equipment with rubber parts.

Administer one dose (2 mL) by intramuscular injection in the neck in the area behind the ear, according to the following schedule:

Vaccination schedule

In the case of low to medium levels of maternally derived antibodies against PCV-2 a single vaccination (2 mL) to pigs from 3 weeks of age.

When higher levels of maternally derived PCV-2 antibodies are expected, the following schedule of two vaccinations is advised: the first injection (2 mL) can be given from an age of 3-5 days, the second injection (2 mL) 2-3 weeks later.

Batch Exp