




**Porcilis<sup>®</sup> PCV**

# **Protection and Profit** *from **START** to **FINISH***



## **Porcilis<sup>®</sup> PCV**

**Long-term PCV2 immunity for performance and profit**

-  A single vaccination (2 mL) to pigs from 3 weeks of age\*
-  Rapid onset of immunity with long-term protection
-  Reduces viral load and shedding
-  Reduces mortality\*
-  Reduces weight loss\*
-  Increases average daily weight gain\*

\* During the fattening period. Refer to registered product label for full claim details.  
Refer to Technical brochure for details of trial data.

Intervet Australia Pty Limited trading as MSD Animal Health ABN 79 008 467 034  
Toll free 1800 033 461 [www.msd-animal-health.com.au](http://www.msd-animal-health.com.au)  
Amanda Vardanega 0427 011 579

 **MSD**  
Animal Health



# Porcilis® PCV

Less virus, more profit

## Porcilis® PCV Vaccine

### ACTIVE CONSTITUENT:

Porcine Circovirus Type 2 ORF2 subunit antigen:  $\geq 4.5 \log_2$  antibody titre per dose of 2 mL.

Adjuvant: 25 mg dl- $\alpha$ -tocopheryl acetate.

346 mg light liquid paraffin.

For active immunisation of pigs to reduce the virus load in blood and lymphoid tissues and to reduce mortality and weight loss associated with Porcine circovirus type 2 related infection (PCVD) occurring during the fattening period.

Onset of immunity: 2 weeks

Duration of immunity: 22 weeks

A single dose breaks through up to medium levels of maternally derived antibodies in piglets.

A double dose breaks through medium to high levels of maternally derived antibodies.

### DIRECTIONS FOR USE:

**SHAKE WELL BEFORE USE. Use the entire contents within 8 hours of opening.**

### RESTRAINTS:

**DO NOT mix with any other vaccine or immunological product.**

### SIDE EFFECTS:

A transient rise in body temperature may occur until 2 days after vaccination. Some piglets may be depressed and show a reduced feed intake for up to 5 days. A transient local reaction at the injection site may be observed in the form of a hard and sometimes warm swelling (up to a diameter of 7 cm) diminishing over a period of approximately 14-21 days. A hypersensitivity-like reaction may occur; the animals

will recover in several minutes without further treatment.

### DOSAGE AND ADMINISTRATION:

The vaccine is a liquid and ready-to-use product. Before using the vaccine allow it to reach room temperature (15-25°C) and shake well before use. Use sterile syringes and needles. Avoid introduction of contamination. Avoid use of vaccination equipment with rubber parts. Administer one dose (2 mL) by intramuscular injection in the neck in the area behind the ear, according to the following schedule:

#### Vaccination Schedule

In the case of low to medium levels of maternally derived antibodies against PCV-2 a single vaccination (2 mL) to pigs from 3 weeks of age. When higher levels of maternally derived PCV-2 antibodies are expected, the following schedule of two vaccinations is advised: the first injection (2 mL) can be given from an age of 3-5 days, the second injection (2 mL) 2-3 weeks later.

### DRUG INTERACTIONS:

Compatibility studies with concurrent use of other veterinary products were not performed.

### WITHHOLDING PERIOD:

**Zero (0) days.**

### TRADE ADVICE: EXPORT

SLAUGHTER INTERVAL (ESI):

ESI not required.

### FIRST AID:

If poisoning occurs contact a doctor or Poisons Information Centre.

*Phone Australia 13 11 26.*

### ADDITIONAL USER SAFETY:

Take care to avoid self-injection. This product contains mineral oil. In the event of self-administration, seek prompt medical attention and take this package leaflet/carton with you. Accidental self-administration may result in local bruising, severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If pain persists for more than 12 hours after medical examination, seek medical advice again.

### To the physician:

This product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert, PROMPT, surgical attention is required and may necessitate early incision of the injected area, especially where there is involvement of finger pulp or tendon.

Additional information is listed in the Material Safety Data Sheet.

**Dispose** of empty container by wrapping with paper and putting in garbage.

**Store** between 2°C and 8°C (Refrigerate. Do not freeze). Protect from light.

### Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

APVMA Approval No.: 64025/47209

MSD Animal Health  
Level 1 - Building A  
26 Talavera Road  
Macquarie Park, NSW 2113

Phone: 1800 033 461

© Registered Trademark

