

FOR ANIMAL TREATMENT ONLY

Nobilis[®] CAV P4 Vaccine

ACTIVE CONSTITUENT: Chicken Anaemia Virus living, attenuated strain 26P4.

Minimum titre at end of shelf life 3.0 log₁₀ TCID₅₀ per dose.

For protection of chicken progeny against Chicken Anaemia Virus (CAV) disease by immunisation of breeders.

DIRECTIONS FOR USE

Dosage and administration
USE ALL PRODUCT WITHIN 4 HOURS
OF RECONSTITUTION.
MIX WELL BY GENTLE SHAKING.

Vaccinate healthy animals only.

For intramuscular (IM) or subcutaneous (SC) injection.

Reconstitute the vaccine in sterile Phosphate Buffered Saline (PBS) Diluent. Select amount of diluent to be used on number of doses per vial (200 mL per 1000 doses). Administer 0.2 mL of the vaccine by IM or SC injection.

General directions

Wash and disinfect hands and equipment after vaccinating.

Vaccination program

The optimum time and method of administration depend largely upon the local situation. Therefore, the advice of a veterinarian should be sought.

The vaccine can be administered to chickens from 6 weeks of age onwards.

To make sure that adequate antibody levels are present at the onset of lay, breeding stock has to be vaccinated at least 6 weeks before this onset.

Immunity

The indication of high neutralising antibody levels is generally dependent on the health and condition of the birds. Hygiene and management are also important in the post vaccination period. Breeding stock have to be vaccinated by at least 6 weeks before onset of lay. If vaccination has been carried out correctly the protective antibody level will be maintained for at least one laying period.

WITHHOLDING PERIODS: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI):
Zero (0) days. Before using this product,
confirm the current ESI from Intervet
Australia Pty Limited on 1800 033 461
or APVMA website
(www.apvma.gov.au/residues).

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

ADDITIONAL USER SAFETY INFORMATION

Take care to avoid self-injection. In the event of self-administration, seek medical attention if you are concerned. Show the package leaflet or label to the Medical Practitioner.

DISPOSAL

Any surplus vaccine should be disposed by burning or boiling. Dispose of vial/container in a designated and appropriately labelled 'biologicals' container.

Discarded needles/sharps should be placed in a designated and appropriately labelled 'sharps' container.

STORAGE

Store below -18°C (deep freeze).

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product

causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

Intervet Australia Pty Limited
(trading as MSD Animal Health)
91-105 Harpin Street,
Bendigo East VIC 3550
Phone: 1800 033 461

APVMA Approval No.: 58408/130888

® Registered trademark

© Copyright 2021

