



Product Name: NOBILIS CAV P4 VACCINE
APVMA Approval No: 58408-130888

Label Name:	NOBILIS CAV P4 VACCINE
Signal Headings:	FOR ANIMAL TREATMENT ONLY
Constituent Statements:	Chicken Anaemia Virus living, attenuated strain 26P4. Minimum titre at end of shelf-life 3.0 log ₁₀ TCID ₅₀ per dose.
Claims:	For protection of chicken progeny against Chicken Anaemia Virus (CAV) disease by immunisation of breeders.
Net Contents:	1 to 25, 50 vials of 500 Doses and 1000 Doses.
Directions for Use:	
Restrictions:	
Contraindications:	
Precautions:	
Side Effects:	
Dosage and Administration:	<p>USE ALL PRODUCT WITHIN 4 HOURS OF RECONSTITUTION. MIX WELL BY GENTLE SHAKING. Vaccinate healthy animals only.</p> <p>For intramuscular (IM) or subcutaneous (SC) injection. Reconstitute the vaccine in sterile Phosphate Buffered Saline (PBS) Diluent. Select amount of diluent to be used on number of doses per vial (200 mL per 1000 doses). Administer 0.2 mL of the vaccine by IM or SC injection.</p>

General Directions:	<p>Wash and disinfect hands and equipment after vaccinating.</p> <p>Vaccination program The optimum time and method of administration depend largely upon the local situation. Therefore, the advice of a veterinarian should be sought. The vaccine can be administered to chickens from 6 weeks of age onwards. To make sure that adequate antibody levels are present at the onset of lay, breeding stock has to be vaccinated at least 6 weeks before this onset.</p> <p>Immunity The indication of high neutralising antibody levels is generally dependent on the health and condition of the birds. Hygiene and management are also important in the post vaccination period. Breeding stock have to be vaccinated by at least 6 weeks before onset of lay. If vaccination has been carried out correctly the protective antibody level will be maintained for at least one laying period.</p>
Withholding Periods:	Zero (0) days.
Trade Advice:	EXPORT SLAUGHTER INTERVAL(ESI): Zero (0) days. Before using this product, confirm the current ESI from Intervet Australia Pty Limited on 1800 033 461 or the APVMA website (www.apvma.gov.au/residues).
Safety Directions:	
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.
First Aid Warnings:	
Additional User Safety:	Take care to avoid self-injection. In the event of self-administration, seek medical attention if you are concerned. Show the package leaflet or label to the Medical Practitioner.
Environmental Statements:	
Disposal:	<p>Any surplus vaccine should be disposed by burning or boiling.</p> <p>Dispose of vial/container in a designated and appropriately labelled biologicals container.</p> <p>Discarded needles/sharps should be placed in a designated and appropriately labelled 'sharps' container.</p>
Storage:	Store below -18°C (deep freeze).