Product Name: PORCILIS APPVAC PLEUROPNEUMONIA VACCINE FOR PIGS

APVMA Approval No: 63910/125814

APVMA	
RLP APPROVED	
	/

Label Name:	PORCILIS APPVAC PLEUROPNEUMONIA VACCINE FOR PIGS
Signal Headings:	FOR ANIMAL TREATMENT ONLY
Constituent Statements:	ACTIVE CONSTITUENTS  Inactivated adjuvanted vaccine containing whole cell Actinobacillus pleuropneumoniae serotypes 1, 7 and 15: Minimum 1x10^8 cells/mL of each antigen including a minimum 18 haemolysin units/mL.  Preservative: 0.1 mg/mL Thiomersal.
Claims:	An aid in the control of pleuropneumonia in pigs, caused by Actinobacillus pleuropneumoniae serotypes 1, 7 and 15.
Net Contents:	1 x 100 mL (50 doses) 1 x 250 mL (125 doses) 1 x 500 mL (250 doses)
Directions for Use:	
Restraints:	
Contraindications:	
Precautions:	
Side Effects:	
Dosage and Administration:	Discard if previously frozen. Shake well to mix before use and keep mixed during use. Use all product within 24 hours of opening.  2 mL by intramuscular injection.

Vaccinate all pigs at around 8 weeks of age and again 2-3 weeks later.

The recommended site of injection is that used for routine injection in pigs i.e. behind and below the ear.

## Equipment

The pack is designed for use with an automatic vaccination gun. For sterilisation of automatic vaccinator see enclosed leaflet.

## CAUTION: AVOID CARCASS DAMAGE

- 1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.
- 2. Maintain cleanliness at all times.
- 3. Keep needles sharp and clean. Replace frequently.
- 4. Use needles of appropriate gauge and length.
- 5. As far as possible avoid injection of pigs that are wet or under dusty conditions.
- 6. This product must be injected only into muscle tissue.
- 7. If possible inject into muscle tissue on side of neck.

## Connecting pack to automatic vaccinator

Connect the tube firmly into the automatic vaccinator. The pack may be suspended by means of a carrying strap. Eject air from the vaccinator and tube by depressing plunger until vaccine flows through needle.

## Resealing vaccine pack

Additional User

Safety:

For best practice vaccine should be used during one administration. If however, unused vaccine is to be used within the following 24 hours, close clip as close as possible to the open end of the tube. Remove tube from the vaccinator gun. Store between 2°C and 8°C (refrigerate, do not freeze). When re-using, cut tube between clip and vaccine pack and reconnect to the vaccinator gun. Discard if previously frozen.

General Directions:	Porcilis APPvac is a killed, whole cell vaccine containing Actinobacillus pleuropneumoniae (serovars 1, 7 and 15). It is recommended as an aid in control of pleuropneumonia in pigs. The adjuvant system increases the level of specific antibodies which act as an aid in protecting pigs through to market age.

	Porcilis APPvac has been shown to be safe when administered to pigs around 8 weeks of age at the recommended dose rate.
Withholding Periods:	WITHHOLDING PERIODS: Zero (0) days.
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.
Safety Directions:	
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	WARNING – Contains formaldehyde.

Take care to avoid self-injection.

	Accidental self-injection may result in local bruising, pain and swelling. In the event of self-injection, seek medical attention and show the package leaflet or the label, to the Medical Practitioner.
Environmental Statements:	
Disposal:	Dispose of empty container by wrapping with paper and putting in garbage
Storage:	Store between 2°C and 8°C (refrigerate, do not freeze). Protect from light. KEEP OUT OF REACH OF CHILDREN.